

PEAK PRECISION NUTRITION



Glut-4 Reg LPT™

ADVANCED BLOOD GLUCOSE & METABOLIC SUPPORT†

Glut-4 Reg LPT™ is a specialized liposomal tablet designed to support blood glucose regulation, insulin sensitivity, and metabolic function. This formula combines clinically studied ingredients to optimize glucose utilization and promote overall metabolic health.



SUPPORTS HEALTHY BLOOD SUGAR LEVELS†

Regulates glucose absorption, enhances insulin sensitivity, and reduces liver glucose production.



**Supports Healthy
Blood Sugar & Insulin Levels†**

PROMOTES METABOLIC HEALTH†

Boosts energy production by increasing glycolysis, supporting thyroid function, and improving mitochondrial efficiency.

ANTIOXIDANT & VASCULAR SUPPORT†

Promotes vascular health by increasing nitric oxide, reducing oxidative stress, and lowering inflammation.

Yin J, et al. Effects and mechanisms of berberine in diabetes treatment. *Acta Pharm Sinica B*. 2012;2(4):327-34.

Lan J, et al. Meta-analysis of the effect and safety of berberine in the treatment of type 2 diabetes mellitus, hyperlipemia, and hypertension. *J Ethnopharmacol*. 2015;161:69-81.

Zhang Y, et al. Treatment of Type 2 diabetes and dyslipidemia with the natural plant alkaloid berberine. *J Clin Endocrinol Metab*. 2008;93(7):2559-65.

Dong H, et al. Berberine in the treatment of type 2 diabetes mellitus: a systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2012;2012:591654.

Kleefstra N, Houweling ST, Bakker SJ, Verhoeven S, Gans RO, Meyboom de Jon B, Bilo HJ. Effect of chromium supplementation on glucose metabolism and lipids: a systematic review of randomized controlled trials. *Diabetes Care* 2007;30:2154-2163.

Lau F, Bagchi M, Sen C, Bagchi D. Nutrigenomic basis of beneficial effects of chromium (III) on obesity and diabetes. *Mol Cell Biochem*. 2008;317:1-10.

Broadhurst CL, Domenico P. Clinical studies on chromium picolinate supplementation in diabetes mellitus—a review. *Diabetes Technol Ther*. 2006;8(6):677-87.

Lau FC, Bagchi M, Sen CK, Bagchi D. Nutrigenomic basis of beneficial effects of chromium(III) on obesity and diabetes. *Mol Cell Biochem*. 2008 Oct;317(1-2):1-10. Epub 2008 Jul 18.

Hou WK, Xian YX, Zhang L, Lai H, Hou XG, Xu YX, Yu T, Xu FY, Song J, Fu CL, Zhang WW, Chen L. Influence of blood glucose on the expression of glucose transporter proteins 1 and 3 in the brain of diabetic rats. *Chin Med J*. 2007;119:1704-9.

Mahmood T, Qureshi IZ, Iqbal MJ. Histopathological and biochemical changes in rat thyroid following acute exposure to hexavalent chromium. *Histol Histopathol*. 2010;25(11):1355-70.

SUPPLEMENT FACTS

Serving Size: 2 Chewable Tablets
Servings Per Container: 30

	Amount Per Serving	% Daily Value
Berberine HCL (Berberis aristate) (Root) (std. 97%)	120 mg	**
Bitter Melon Extract (Momordica Charantia Linn.) (Fruit) (std. to 10% Charantins)	50 mg	**
Chromium (as chromium GTF)	800 mcg	2,286%
Micro PQQ	20 mg	**
Vanadium (as Vanadyl sulphate)	4 mg	**
Vitamin B6 (as Pyridoxal-5-Phosphate)	10 mg	588%
Liposomal Blend (LPT™)	1600 mg	**
Dicalcium Phosphate, Dextrose, Sunflower Lecithin, Beta Cyclodextrin		

** Daily Value not established.

Other Ingredients:

Silicon Dioxide, Magnesium Stearate.

Contains no salt, soy, yeast, dairy, gluten, artificial colors, flavors, or preservatives.

THIS PRODUCT IS GLUTEN FREE.

Tamper Resistant Seal. If Seal Is Broken, Do Not Use.

If you are pregnant, nursing or taking any medications, consult with your healthcare provider before use.

Keep out of the reach of children.

RECOMMENDED USE

Swallow one (1) tablet with water, twice daily, or as directed by a healthcare professional. **Do not chew.**

SCIENCE BEHIND LPT™

LIPOTABS™ (LPT™) are safe, chewable, oral liposomal tablets manufactured using patent pending, revolutionary small molecule technology in order to support improved bioavailability and effects on the body.[†]

Misra HS, Raipurohit YS, Khairnar NP. Pyrroloquinoline-quinone and its versatile roles in biological processes. J Biosci. 2012;37(2):313-25.

Takada M, Sumi M, Maeda A, et al. Pyrroloquinoline quinone, a novel protein tyrosine phosphatase 1B inhibitor, activates insulin signaling in C2C12 myotubes and improves impaired glucose tolerance in diabetic KK-A(y) mice. Biochem Biophys Res Commun. 2012;428(2):315-20.

McInerney MF, Seidel MJ, Nguyen JM, et al. Effects of a 33 residue interleukin-1 beta peptide and the antioxidant PQQ on interleukin-1 beta-mediated inhibition of glucose-stimulated insulin release from cultured mouse pancreatic islets. Res Commun Mol Pathol Pharmacol. 1996;94(2):115-28.

Nix WA, et al. Vitamin B status in patients with type 2 diabetes with and without incipient nephropathy. Diabetes Res Clin Pract. 2015;107(1):157-65.

Khobrani M, et al. Impact of vitamin B6 deficiency on the severity of diabetic peripheral neuropathy – a cross-sectional study. Saudi Pharm J. 2023;31(5):655-58.

Smith DM, Pickering RM, Lewith GT. A systematic review of vanadium oral supplements for glycaemic control in type 2 diabetes mellitus. QJM. 2008;101(5):351-8.

[No Authors Listed]. Vanadium (vanadyl sulfate). Monograph. Altern Med Rev. 2009;14(2):177-80.

[†] These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



Copyright © 2025 James B. LaValle, Natural Formulations All rights reserved.

No part of this material may be used or reproduced in any manner whatsoever, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the author.

This material is provided for educational and informational purposes only to licensed health care professionals. This information is obtained from sources believed to be reliable, but its accuracy cannot be guaranteed. Herbs and other natural substances are very powerful and can occasionally cause dangerous allergic reactions in a small percentage of the population. Licensed health care professionals should rely on sound professional judgment when recommending herbs and natural medicines to specific individuals. Individual use of herbs and natural medicines should be supervised by an appropriate health care professional. The use of any specific product should always be in accordance with the manufacturer's directions.